



Corona (COVID-19) time musings: Our involvement in COVID-19 pathogenesis, diagnosis, treatment and vaccine planning

1. Pathogenesis

The 2020 coronavirus epidemic is characterized by high infectious rates and relatively high mortality, especially among the elderly over the age of 80. In this issue we published two reports related to the COVID-19 infection [1,2]. In most of those severe cases of coronavirus infection, the clinical expression is characterized not only by a fever, cough and other constitutional symptoms, but also by a clinical constellation including a cytokine storm, respiratory failure and eventually death [3]. Among the aberrant laboratory findings which characterize these clinical expressions, one can find leukopenia, increased liver function tests and ferritin levels reaching the hundreds and sometimes thousands of units. This panoply of findings reminds us of four clinical conditions that we have described in the past under the new syndrome — hyperferritinemic syndrome [4–6]. These four conditions include macrophage activating syndrome (MAS) [4], catastrophic antiphospholipid syndrome (cAPS) [7], septic shock and an adverse reaction to the biological compound anti-CD-28 [8]. This reaction to the drug was observed in a short therapeutic trial published in the *New England Journal of Medicine* in which 6 individuals were hospitalized in the intensive care unit with a cytokine storm following the therapy [8]. All the above conditions are characterized by high ferritin levels, a cytokine storm (especially of inflammatory cytokines such as IL-1, IL-6, IL-17, etc.) and a high mortality rate that is over 50%. The 6 hospitalized individuals from the aforementioned anti-CD-28 trial did not die as they were immediately rushed to the intensive care unit [8,9]. In the other 3 conditions, we were able to reduce the mortality rate significantly below 50% using combined therapies with corticosteroids, antibiotics, IVIG (intra-venous gamma-globulins) and anticytokine therapy (anti-IL-1 and/or anti-IL-6) [10–12]. The resemblance between these clinical conditions and those of COVID-19, suggests that the very severe cases of the coronavirus are associated with a clinical picture similar to that of the macrophage activating syndrome (MAS), which is unsurprisingly associated with high levels of ferritin. Therefore, the enigma regarding the high mortality rate associated with COVID-19 is most probably explained by a cytokine storm. Understanding this pathogenetic pathway may lead to better therapy and better survival rates. We will consider this further in our discussion of therapy.

2. Diagnosis (Ferritin, sCD163)

It goes without saying that the diagnosis of a COVID-19 infection is based on the presence of the virus in oro-pharyngeal swabs taken at the right time, as well as on the presence of IgM (and eventually IgG) antibodies against the virus. In severe cases, the clinical picture diagnosis

may be supported by the clinical manifestations of severe respiratory failure syndrome, as well as by blood tests indicating leukopenia, increased liver function tests and high levels of ferritin [13]. Additionally, we would also like to suggest to test for soluble CD-163 (sCD-163), which represents the activation of macrophages [14,15], the level of which was found to increase in MAS and to parallel the ferritin level [15].

In the past, we have shown this parameter to be involved in parallel to the ferritin being increased during the acute stage of the inflammation (14, 15). A commercial kit for analysing the sCD-163 is available in the market. The question of the pathogenesis relates to whether the ferritin is just an epiphenomenon that can be used for diagnostic purposes, or is the ferritin involved in the vicious cycle perpetuating and exasperating the inflammation. Recently together with P. Ruscitti et al. [16], we have found the role of the H-chain of the ferritin in activating macrophages to increase the secretion of inflammatory cytokines. Thus the circle is completed on our understanding of the pathogenesis of the hyperferritinemic syndrome including the infection with Covid-19.

3. Therapy (IVIG, anti-IL-1, anti-IL-6)

Due to the very rapid distribution of scientific knowledge today, several optional therapies have been already suggested in a small study scales to be beneficial such as the application of chloroquine (plaquenil), remdesivir and others [3]. However, understanding the pathogenesis of the virus causing the cytokine storm one would recommend the addition of anticytokine biological agents. Such as anti-IL-1 (anakinra) or anti-IL-6 (tocilizumab (TCZ)). These drugs are available especially in rheumatological autoimmune inflammatory conditions and have very good therapeutic results. But I would like to refer to an additional suggested therapy, namely: IVIG. IVIG which is composed of immunoglobulin extracted from 20,000 healthy normal subjects was reported in the past to be effective in several cases of MAS as well as in septic shock. Previously we have reported that IVIG *per se* contains also panoply of anti-viral antibodies [17]. This knowledge has been used in another epidemic in the past of the West Nile Fever (WNF) in New York, a geographical area that by and large is not exposed to this virus. An IVIG that was generated in Israel - an endemic area for the West Nile Fever virus and extracted from healthy Israeli convalescence blood donors was found to be effective in reducing the morbidity of the West Nile Fever in New York [18–20]. IVIG *per se* can be helpful in an infectious situation by the transfer of a normal innate immune system of healthy subject to the infected individual [21,22]. Having almost no side effects [23], it would be recommended to deliver the IVIG in a dose of 2 g per Kg bodyweight in 4 days (according to the bodyweight of the

person). Needless to say, that an IVIG produced from a large number of SERA derived from convalescence subjects from the viral infection will be much more effective and could be delivered in smaller doses. Preliminary studies have already been performed in several sites in the world [13]. We would urge to produce what we have called in the past specific IVIG (sIVIG) in which this specific convalescence IgG would be extracted and could be used really in minimal amounts [24–27]. There is a possibility to affinity purify this IVIG either on a column constructed from peptides of the spikes of the virus or from peptides known to be the constituents of the virus as was delineated by our recent paper [28].

4. Vaccine (Avoid side effects)

We all are expecting the vaccine production trials to materialise quickly. We believe that once the vaccine found to be effective (most probably on a theoretical basis) it will be distributed to millions or billions of people. We believe that this vaccine will be approved through an expedited process thus not necessarily enabling surveillance due to the shortness of time thus eventual side effects of the vaccine could not be evaluated. The amino acid sequences of the virus like in other viruses, might have a cross-reaction with the human body sequences [28–31]. Therefore, one of the side effects of giving a MASS vaccine could be an emergence of autoimmune diseases especially in individuals who are genetically prone for autoimmunity [28–31]. Actually the coronavirus was reported to induce retinal autoimmune disease in an experimental model [32]. We have recently delineated the amino acid penta-peptides of the virus and selected those who are immunogenic yet have no similarities to the human constituents and we believe that any vaccine which can be produced by such a method will reduce significantly the eventual side effect of induction of autoimmune diseases [28]. Those immunogenic peptides can also be used to generate an affinity column to extract the specific antibodies to the virus existing in the SERA of convalescence subjects from covid-19. (See section Therapy).

5. Conclusion

Better understanding of the pathogenesis of the infection with the covid-19 which in selected cases may lead to a similar clinical picture of macrophage activating syndrome (MAS) with its associated cytokine storm may bring to an improved diagnostic measurements. Such precision medicine may help in early diagnosis of deterioration into the severe clinical conditions. Moreover, understanding this pathogenesis may lead to a better therapeutic measurements which may entail also anti cytokine therapy as well as adding to the therapeutic regimen IVIG. If possible harnessing specific IVIG which is enriched with anti COVID-19 antibodies extracted from the SERA of patients who recovered from the viral infection.

Acknowledgement

This work is supported by the grant of the Government of the Russian Federation for the state support of scientific research carried out under the supervision of leading scientists, agreement 14.W03.31.0009.

References

- [1] McGonagle Dennis, Sharif Kassem, O'Regan Anthony, Bridgewood Charlie. Interleukin-6 use in COVID-19 pneumonia related macrophage activation syndrome. *Autoimmun Rev* 2020. [DOI10.1056/NEJM.e2005477](#). [epub.ahead of print].
- [2] Marotto Daniela, Sarzi-Puttini Piercarlo. What is the role of rheumatologists in the era of COVID-19? *Autoimmunity Rev* 2020. *Autoimmunity reviews* this issue.
- [3] Baden Lindsey R, Rubin Eric J. Covid-19 – The search for effective therapy. Editorial. *N Engl J Med* 2020. Published on March 18, 2020, [DOI 10.1056/NEJM](#). [epub.ahead of print].
- [4] Rosario C, Zandman-Goddard G, Meyron-Holtz EG, D'Cruz DP, Shoenfeld Y. The Hyperferritinemic syndrome: macrophage activation syndrome, Still's disease, septic shock and catastrophic antiphospholipid syndrome. *BMC Med* 2013;11:185–96.
- [5] Zandman-Goddard G, Orbach H, Agmon-Levin N, Boaz M, Amital H, Szekancz Z, et al. Hyperferritinemia is associated with serologic antiphospholipid syndrome in SLE patients. *Clin Rev Allergy Immunol* 2013;44:23–30.
- [6] Zandman-Goddard G, Shoenfeld Y. Editorial: Hemophagocytic Syndrome with hyperferritinemia: a stormy immunological response. *IMAJ* 2013;15:187–8. (This is an Editorial for article: Decruz D. et al. N2010 Adult-Onset Still's Disease complicated by hemophagocytic syndrome and catastrophic antiphospholipid syndrome resulting in four limb amputation. *IMAJ* 2013; 15: 192–194).
- [7] Agmon-Levin N, Rosario C, Porat-Katz B-S, Zandman-Goddard G, Meroni P, Cervera R, et al. Ferritin in the antiphospholipid syndrome and its catastrophic variant (cAPS). *Lupus* 2013;22:1327–35.
- [8] Suntharalingam G, Perry MR, Ward S, Brett SJ, Castello-Cortes A, Brunner MD, et al. Cytokine storm in a phase 1 trial of the anti-CD28 monoclonal antibody TGN1412. *N Engl J Med* 2006;355(10):1018–28.
- [9] Guo Xi-Zhi J, Thomas Paul G. New fronts emerge in the influenza cytokine storm. *Semin Immunopathol* 2017;39:541–50.
- [10] Cervera R, Rodríguez-Pintó I, Colafrancesco S, Conti F, Valesini G, Rosário C, et al. 14th international congress on antiphospholipid antibodies task force report on catastrophic antiphospholipid syndrome. *Autoimmun Rev* 2014;13:699–707.
- [11] Rodríguez-Pintó Ignasi, Moitinho Marta, Santacreu Irene, Shoenfeld Yehuda, Erkan Doruk, Espinosa Gerard, et al. Catastrophic antiphospholipid syndrome (CAPS): Descriptive analysis of 500 patients from the International CAPS Registry. *Autoimmun Rev* 2016;15:1120–4.
- [12] Carmi O, Berla M, Shoenfeld Y, Levy Y. Diagnosis and management of catastrophic antiphospholipid syndrome. *Expert Rev Hematol* 2017;10(4):365–74.
- [13] Mehta Puja, McAuley Daniel F, Brown Michael, Sanchez Emilie, Tattersall Rachel S, Manson Jessica J, et al. UK. COVID-19: consider cytokine storm syndromes and immunosuppression. [www.thelancet.com](#) Published online March 13 2020. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](#).
- [14] Colafrancesco S, Priori R, Alessandri C, Astorri E, Perricone C, Blank M, et al. The Hyperferritinemic syndromes and CD163: a marker of macrophage activation. *IMAJ* 2014;16:662–3.
- [15] Colafrancesco S, Priori R, Alessandri C, Astorri E, Perricone C, Blank M, et al. sCD163 in AOSD: a biomarker for macrophage activation related to hyperferritinemia. *Immunol Res* 2014;60:177–83.
- [16] Ruscitti Piero, Berardicurti Onorina, Cipriani Paolo, Iagnocco Annamaria, Shoenfeld Yehuda. Severe hyper-inflammatory COVID-19, another piece in the puzzle of the "hyperferritinemic syndrome". *Rheumatol Point View* 2020. (Submitted for publication).
- [17] Krause I, Wu R, Sherer Y, Patanik M, Peter J, Shoenfeld Y. In vitro antiviral and antibacterial activity of commercial intravenous immunoglobulin preparations - a potential role for adjuvant intravenous immunoglobulin therapy in infectious diseases. *Transfus Med* 2002;12:133–9.
- [18] Planitzer CB, Mofro J, Kreil TR. West Nile virus neutralization by US plasma-derived immunoglobulin products. *J Infect Dis* 2007;196(3):435–40.
- [19] Haley M, Retter AS, Fowler D, Gea-Banacloche J, O'Grady NP. The role for intravenous immunoglobulin in the treatment of West Nile virus encephalitis. *Clin Infect Dis* 2003;37:e88–90.
- [20] Shomoni Z, Niven MJ, Pitlick S, Bulvik S. Treatment of West Nile virus encephalitis with intravenous immunoglobulin. *Emerg Infect Dis* 2001;7(4):759.
- [21] Domizi R, Adrario E, Damiani E, Scorsella C, Carsetti A, Giaccaglia P, et al. IgM-enriched immunoglobulins (pentaglobin) may improve the microcirculation in sepsis: a pilot randomized trial. *Ann Intensive Care* 2019;9(1):135. [https://doi.org/10.1186/s13613-019-0609-5](#).
- [22] Yang Y, Yu X, Zhang F, Zia Y. Evaluation of the effect of intravenous immunoglobulin dosing n mortality in patients with sepsis: A network meta-analysis. *Clin Ther* 2019;41(9). (1838.e4).
- [23] Orbach H, Katz U, Sherer Y, Shoenfeld Y. Intravenous immunoglobulin (IVIG) adverse effects and safe administration. *Clin Rev Allergy Immunol* 2005;29:173–84.
- [24] Blank M, Nur I, Toub O, Aron-Maor A, Shoenfeld Y. Toward molecular targeting with specific intravenous immunoglobulin preparation. *Clin Rev Allergy Immunol* 2005;29:213–7.
- [25] Svetlicky N, Ortega-Hernandez AD, Mouthon L, Guillemin L, Thiesen HJ, Altman A, et al. The advantage of specific intravenous immunoglobulin (sIVIG) on regular IVIG: Experience of the last decade. *J Clin Immunol* 2013;33(Suppl. 1):S27–32.
- [26] Blank M, Bashi T, Shoenfeld Y. Idiotype-specific intravenous immunoglobulin (IVIG) for therapy of autoimmune diseases. Chapter 18. In: Steinitz Michael, editor. *Human Monoclonal Antibodies: Methods and Protocols*. Methods in Molecular Biology, 1060. 2014. p. 353–61.
- [27] Svetlicky N, Kivity S, Odeh Q, Shovman O, Gertel S, Amital H, et al. Anti-citrullinated protein antibody-specific intravenous immunoglobulin attenuates collagen-induced arthritis in mice. *Clin Exp Immunol* 2015;182:241–50. [https://doi.org/10.1111/cei.12673](#).
- [28] Kanduc D, Shoenfeld Y. On the molecular determinants and the mechanism of the SARS-CoV-2 attack. 2020. (Submitted).
- [29] Darja Kanduc, Yehuda Shoenfeld. Inter-pathogen peptide sharing and the original antigenic sin: solving a paradox. *Open Immunol J* 2018;08:16–27 Content list available at: [www.benthamopen.com/TOLJ/](#) [https://doi.org/10.2174/1874226201808010016](#).
- [30] Kanduc D, Shoenfeld Y. Human papillomavirus epitope mimicry and autoimmunity: The molecular truth of peptide sharing. *Pathobiology* 2019. [https://doi.org/10.1159/000502889](#). Published online.

- [31] Darja Kanduc, Yehuda Shoenfeld. From anti-EBV immune responses to the EBV disease via cross-reactivity. *Infection International* 2019;7(5):1–21. (Received September 01, 2019; accepted September 10, 2019).
- [32] Detrick Barbara, Hooks John J. Robert Nussenblatt. Infections associated with retinal autoimmunity. Chapter No.57, pp: 943–957. In: Shoenfeld Yehuda, Agmon-Levin Nancy, Rose Noel R, editors. BOOK: “Infection and Autoimmunity”. Elsevier BV Academic Press Publication; 2015. p. 1–1036.

Yehuda Shoenfeld^{a,b,*}

^a *Zabludowicz Center for Autoimmune Diseases, Sheba Medical Center, affiliated to Tel-Aviv University, Israel*

^b *Laboratory of the Mosaics of Autoimmunity, Saint Petersburg State University, Saint-Petersburg, Russian Federation*

E-mail address: Yehuda.Shoenfeld@sheba.health.gov.il.

* Zabludowicz Center for Autoimmune Diseases, Sheba Medical Center, affiliated to Tel-Aviv University, Israel.